

Dark torsion as the cosmic speed-up

Gabriel R. Bengochea^{1,2,*} and Rafael Ferraro^{1,2,†}

¹*Instituto de Astronomía y Física del Espacio, Casilla de Correo 67, Sucursal 28, 1428 Buenos Aires, Argentina*

²*Departamento de Física, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Ciudad Universitaria, Pabellón I, 1428 Buenos Aires, Argentina*

It is shown that the recently detected acceleration of the universe can be understood by considering a modification of the teleparallel equivalent of General Relativity (TEGR), with no need of dark energy. The solution also exhibits phases dominated by matter and radiation as expected in the standard cosmological evolution. We perform a joint analysis with measurements of the most recent type Ia supernovae (SNe Ia), Baryon Acoustic Oscillation (BAO) peak and estimates of the CMB shift parameter data to constraint the only new parameter this theory has.

I. INTRODUCTION

The discovery of an unexpected diminution in the observed energy fluxes coming from type Ia supernovae [1, 2] has opened one of the most puzzling and deepest problems in cosmology today. These observations have been interpreted as solid evidence for an accelerating universe dominated by something called dark energy. Although the cosmological constant seems to be the simplest explanation for the phenomenon, several dynamical scenarios have been tried out since 1998 (see e.g., [3, 4, 5]). While some authors sustain the idea of the existence of a dark energy, others propose modifications of the Einstein-Hilbert Lagrangian known as $f(R)$ ([6, 7, 8, 9, 10, 11, 12, 13, 14] or [15, 16, 17] for recent reviews) as a way to obtain a late accelerating expansion. A great difficulty these theories have, from the point of view of the metric formalism, is that the resulting field equations are 4th order equations, which in many cases makes these hard to analyze. Besides, the simplest cases of the kind $f(R) = R - \beta/R^n$ have shown difficulties with, weak field tests [18, 19], gravitational instabilities [20] and do not present a matter dominated era previous to the acceleration era [21, 22]. Alternatively, the Palatini variational approach for such $f(R)$ theories leads to 2nd order field equations, and some authors have achieved to put observational constraints to these theories [23, 24, 25]. However in many cases the equations are still hard to work with, as evidenced by the functional form of the modified Friedmann equation for a generic $f(R)$. Recently, models based on modified teleparallel gravity were presented as an alternative to inflationary models [26, 27]. In this paper we show a cosmological solution for the acceleration of the universe by means of a sort of theories of modified gravity, namely $f(L_T)$, based on a modification of the teleparallel equivalent of General Relativity (TEGR) Lagrangian [28, 29] where the torsion will be the responsible of the observed

acceleration of the universe, and the field equations will always be 2nd order equations.

II. GENERAL CONSIDERATIONS. FIELD EQUATIONS

Teleparallelism [28, 29] uses as dynamical object a vierbein field $\mathbf{e}_i(x^\mu)$, $i = 0, 1, 2, 3$, which is an orthonormal basis for the tangent space at each point x^μ of the manifold: $\mathbf{e}_i \cdot \mathbf{e}_j = \eta_{ij}$, where $\eta_{ij} = \text{diag}(1, -1, -1, -1)$. Each vector \mathbf{e}_i can be described by its components e_i^μ , $\mu = 0, 1, 2, 3$ in a coordinate basis; i.e. $\mathbf{e}_i = e_i^\mu \partial_\mu$. Notice that latin indexes refer to the tangent space, while greek indexes label coordinates on the manifold. The metric tensor is obtained from the dual vierbein as $g_{\mu\nu}(x) = \eta_{ij} e_i^\mu(x) e_j^\nu(x)$. Differing from General Relativity, which uses the torsionless Levi-Civita connection, Teleparallelism uses the curvatureless Weitzenböck connection [30], whose non-null torsion is

$$T_{\mu\nu}^\lambda = \overset{\mathbf{w}}{\Gamma}_{\nu\mu}^\lambda - \overset{\mathbf{w}}{\Gamma}_{\mu\nu}^\lambda = e_i^\lambda (\partial_\mu e_\nu^i - \partial_\nu e_\mu^i) \quad (1)$$

This tensor encompasses all the information about the gravitational field. The TEGR Lagrangian is built with the torsion (1), and its dynamical equations for the vierbein imply the Einstein equations for the metric. The teleparallel Lagrangian is [29, 31, 32],

$$L_T = S_\rho^{\mu\nu} T_{\mu\nu}^\rho \quad (2)$$

where:

$$S_\rho^{\mu\nu} = \frac{1}{2} \left(K^{\mu\nu}_\rho + \delta_\rho^\mu T^{\theta\nu}_\theta - \delta_\rho^\nu T^{\theta\mu}_\theta \right) \quad (3)$$

and $K^{\mu\nu}_\rho$ is the contorsion tensor:

$$K^{\mu\nu}_\rho = -\frac{1}{2} \left(T^{\mu\nu}_\rho - T^{\nu\mu}_\rho - T_\rho^{\mu\nu} \right) \quad (4)$$

which equals the difference between Weitzenböck and Levi-Civita connections.

In this work the gravitational field will be driven by a Lagrangian density that is a function of L_T . Thus the action reads

$$I = \frac{1}{16\pi G} \int d^4x e f(L_T) \quad (5)$$

*Electronic address: gabriel@iafe.uba.ar; CONICET Fellow

†Electronic address: ferraro@iafe.uba.ar; Member of Carrera del Investigador Científico (CONICET, Argentina)

where $e = \det(e_\mu^i) = \sqrt{-g}$. The case $f(L_T) = L_T$ corresponds to TEGR. If matter couples to the metric in the standard form then the variation of the action with respect to the vierbein leads to the equations

$$\begin{aligned} e^{-1} \partial_\mu (e S_i^{\mu\nu}) f'(L_T) - e_i^\lambda T^\rho_{\mu\lambda} S_\rho^{\nu\mu} f'(L_T) + \\ + S_i^{\mu\nu} \partial_\mu (L_T) f''(L_T) + \frac{1}{4} e_i^\nu f(L_T) = \\ = 4\pi G e_i^\rho T_\rho^\nu \end{aligned} \quad (6)$$

where a prime denotes differentiation with respect to L_T , $S_i^{\mu\nu} = e_i^\rho S_\rho^{\mu\nu}$ and $T_{\mu\nu}$ is the matter energy-momentum tensor. The fact that equations (6) are 2nd order makes them simpler than the dynamical equations resulting in $f(R)$ theories.

III. COSMOLOGICAL SOLUTION AND OBSERVATIONAL CONSTRAINTS

We will assume a flat homogeneous and isotropic FRW universe, so

$$e_\mu^i = \text{diag}(1, a(t), a(t), a(t)) \quad (7)$$

where $a(t)$ is the cosmological scale factor. By replacing in (1), (3) and (4) one obtains

$$L_T = S^{\rho\mu\nu} T_{\rho\mu\nu} = -6 \frac{\dot{a}^2}{a^2} = -6 H^2 \quad (8)$$

H being the Hubble parameter $H = \dot{a} a^{-1}$. As a remarkable feature, the scale factor enters the invariant L_T just through the Hubble parameter. The substitution of the vierbein (7) in (6) for $i = 0 = \nu$ yields

$$12 H^2 f'(L_T) + f(L_T) = 16\pi G \rho \quad (9)$$

Besides, the equation $i = 1 = \nu$ is

$$48 H^2 f''(L_T) \dot{H} - f'(L_T) [12 H^2 + 4 \dot{H}] - f(L_T) = 16\pi G p \quad (10)$$

In Eqs. (9-10), $\rho(t)$ and $p(t)$ are the total density and pressure respectively. It can be easily derived that they accomplish the conservation equation

$$\frac{d}{dt}(a^3 \rho) = -3 a^3 H p \quad (11)$$

whatever $f(L_T)$ is. Thus, if the state equation is $p = w\rho$ then ρ evolves as $\rho \propto (1+z)^{3(1+w)}$ (z is the cosmological redshift).

We are interested in obtaining an accelerated expansion without dark energy but driven by torsion. For this we will try with a kind of $f(L_T)$ theories:

$$f(L_T) = L_T - \frac{\alpha}{(-L_T)^n} \quad (12)$$

being α and n real constants to be determined by observational constraints. Although the functional form of

(12) is similar to those considered in $f(R)$ literature, now the guideline towards modified gravity is H instead of R . This fact gives to these theories another interesting feature because H is the most important cosmological variable. For later times the term $-\alpha/(-L_T)^n$ is dominant, while in early times, when $H \rightarrow \infty$, General Relativity is recovered. From (9) along with (12), the modified Friedmann equation results to be

$$H^2 - \frac{(2n+1)\alpha}{6^{n+1}H^{2n}} = \frac{8}{3}\pi G \rho \quad (13)$$

(a functional dependence similar to the results other authors arrived, through different theoretical motivations such as [33, 34]).

Now, replacing $\rho = \rho_{mo}(1+z)^3 + \rho_{ro}(1+z)^4$, and calling $\Omega_i = 8\pi G \rho_{io}/(3H_o^2)$ the contributions of matter and radiation to the total energy density today, Eq. (13) becomes

$$y^n(y-B) = C \quad (14)$$

where $y = H^2/H_o^2$, $B = \Omega_m(1+z)^3 + \Omega_r(1+z)^4$ and $C = \alpha(2n+1)/(6H_o^2)^{-(n+1)}$. The evaluation of this equation for $z = 0$ allows to rephrase the constant C as a function of Ω_i and n : $C = 1 - \Omega_m - \Omega_r$. For $\alpha = 0$ (then $1 = \Omega_m + \Omega_r$) the GR spatially flat Friedmann equation $H^2 = H_o^2 B$ is retrieved. The case $n = 0$ recovers the GR dynamics with cosmological constant $\Omega_\Lambda = 1 - \Omega_m - \Omega_r$. Notice the functional simplicity of (13) compared with its analogue in $f(R)$ theories. Compared with GR, n is the sole new free parameter in (14), since specifying the value of n and Ω_m (Ω_r) the value of α (in units of $H^{2(n+1)}$) is automatically fixed through the relation (13). In order to obtain $H(z)$ we solve numerically the equation (14).

Since the most solid evidence for the acceleration of the universe comes from measurements of luminosity distances for type Ia supernovae, we will use the most recent compilation of 307 SNe Ia events (the *Union* sample) [2], to put constraints in the $n-\Omega_m$ plane. The predicted distance modulus for a supernova at redshift z , for a given set of parameters $\mathbf{P}=(n, \Omega_m)$, is

$$\mu(z | \mathbf{P}) = m - M = 5 \log(d_L) + 25 \quad (15)$$

where m and M are the apparent and absolute magnitudes respectively, and d_L stands for the luminosity distance (in units of megaparsecs),

$$d_L(z; \mathbf{P}) = (1+z) \int_0^z \frac{dz'}{H(z', \mathbf{P})} \quad (16)$$

where $H(z; \mathbf{P})$ is given by the numerical solution of (14). We use a χ^2 statistic to find the best-fit for a set of parameters \mathbf{P} (marginalizing over H_o),

$$\chi_{SNe}^2 = \sum_{i=1}^{N=307} \frac{[\mu_i(z | \mathbf{P}) - \mu_i^{obs}(z)]^2}{\sigma_i^2} \quad (17)$$

where $\mu_i(z | \mathbf{P})$ is defined by (15), μ_i^{obs} and σ_i are the distance modulus and its uncertainty for each observed

value [2]. As it is known, the measurements of SNe Ia are not enough to constrain Ω_m thoroughly. To perform the statistic we also consider, on one hand, the information coming from the BAO (Baryon Acoustic Oscillation) peak detected in the correlation function of luminous red galaxies (LRG) in the Sloan Digital Sky Survey [35]. The observed scale of the peak effectively constraints the quantity (assumed a Λ CDM model),

$$A_{0.35} = D_V(0.35) \frac{\sqrt{\Omega_m H_o^2}}{0.35} = 0.469 \pm 0.017 \quad (18)$$

where $z = 0.35$ is the typical redshift of the LRG and D_V is defined as

$$D_V(z) = \left[\frac{z}{H(z)} \left(\int_0^z \frac{dz'}{H(z')} \right)^2 \right]^{1/3} \quad (19)$$

On the other hand, we have also included in the statistic the CMB shift parameter, which relates the angular diameter distance to the last scattering surface with the angular scale of the first acoustic peak in the CMB power spectrum. In order to do this, we have considered a radiation component $\Omega_r = 5 \times 10^{-5}$. The CMB shift parameter is given by [36],

$$R_{1089} = \sqrt{\Omega_m H_o^2} \int_0^{1089} \frac{dz}{H(z)} = 1.710 \pm 0.019 \quad (20)$$

We can use both parameters since our model presents matter domination at the decoupling time. Figure 1 shows the Hubble diagram for the 307 SNe Ia belonging to the *Union* sample. The curves represent models with values of Ω_m and n obtained from minimizing χ^2 using only SNIa and SNIa+BAO+CMB as well. It is also shown as a reference the Λ CDM model with $\Omega_m = 0.26$. The obtained values for the best-fit to the SNe Ia data only are $\Omega_m = 0.42$ and $n = 1.30$ with the reduced $\chi^2_{\nu} \equiv \chi^2_{min}/\nu \simeq 1.02$ (or equivalently, $\Delta\chi^2_{min} = -1.1$) where ν is the number of degrees of freedom.

In Figure 2 we plot the distance modulus residual ($\Delta\mu$) from the Λ CDM model to better appreciate the discrepancies between our model and Λ CDM.

Figure 3 shows the confidence intervals at 68.3%, 95.4% and 99.7% for the joint probability of the parameters n and Ω_m , having combined the SNe Ia data with BAO and CMB parameters. This analysis yields that the best-fit to all data is achieved with $n = -0.10$ and $\Omega_m = 0.27$ (with a $\chi^2_{min}/\nu \simeq 1.01$, $\Delta\chi^2_{min} = -1.2$) and also the values of the parameters lie in the ranges (at 68.4% c.l.): $n \in [-0.23, 0.03]$; $\Omega_m \in [0.25, 0.29]$.

For our model we have analyzed as well the total and effective equations of state as a function of z . From (13) and (10) along with (12), one can define a torsion contri-




FIG. 1: Hubble diagram for 307 SNe Ia from the *Union* sample [2]. The curves correspond to the concordance model Λ CDM with $\Omega_\Lambda=0.74$ and $\Omega_m=0.26$ (dotted line), and our models with the values corresponding to the best-fit $\Omega_m=0.42$ and $n=1.30$ (dashed line) and also with those coming from the joint analysis of SNIa+BAO+CMB, $\Omega_m=0.27$ and $n=-0.10$ (solid line).




FIG. 2: Distance modulus residual from the Λ CDM model for the same values from Figure 1.

bution to the density and pressure as

$$\begin{aligned} \rho_T &= \frac{3}{8\pi G} \frac{(2n+1)\alpha}{6^{n+1}H^{2n}} \\ p_T &= \frac{\alpha}{8\pi G} \left[(6H^2)^{-(n+1)} \dot{H} [4n(n+1) - 2n] + \right. \\ &\quad \left. - 6n(6H^2)^{-(n+1)} H^2 - \frac{(6H^2)^{-n}}{2} \right] \end{aligned} \quad (21)$$

to rewrite the dynamical equations as

$$H^2 = \frac{8\pi G}{3} (\rho + \rho_T) \quad (22)$$

$$\frac{\ddot{a}}{a} = -\frac{8\pi G}{6} [\rho + \rho_T + 3(p + p_T)] \quad (23)$$

Then, by using (22) and (23) the total and effective equa-




FIG. 3: Confidence intervals at 68.3%, 95.4% y 99.7% in the $n - \Omega_m$ plane coming from combining SNe Ia, BAO and CMB data. The best-fit to this joint analysis is reached with the values $n = -0.10$ and $\Omega_m = 0.27$.

tions of state are written as,

$$w_{tot} \equiv \frac{p + p_T}{\rho + \rho_T} = -1 + \frac{2(1+z)}{3H} \frac{dH}{dz} \quad (24)$$

$$w_{eff} = \frac{p_T}{\rho_T} \quad (25)$$

Figure 4 shows the evolution of the total equation of state w_{tot} as a function of z for our model with the values of the best-fit. There can be observed the last three phases of the evolution of the universe: radiation dominated ($w = 1/3$), matter dominated ($w = 0$) and late acceleration ($w \simeq -1$). Figure 5 shows the effective equation of state coming from the dark torsion contribution. Finally, analyzing (23) we find that our model with $n = -0.10$ predicts that the transition from deceleration to acceleration occurs at $z_{acc} \simeq 0.74$ in good agreement with recent works [37].

An interesting point to be highlighted is that equation (13) reveals that a value of $n > 0$, as the one obtained by considering only SNIa data, implies that the effective dark torsion is of the phantom type [38]. That is, since H decreases toward the present time, the dark torsion density increases instead of diluting with expansion ($w_{eff} < -1$). However, when combining the complete data with SNIa+BAO+CMB we can see from Fig. 3 that it is slightly favored (1σ c.l.) a model with $n \leq 0$.

IV. CONCLUSIONS

A theory $f(L_T)$ based on a modification of the teleparallel equivalent of General Relativity (TEGR) -where torsion is the geometric object describing gravity instead of curvature and its equations are always of 2nd order- is remarkably simpler than $f(R)$ theories. We have tested




FIG. 4: The curves correspond to the total equation of state as a function of z expected for the standard concordance model Λ CDM with $\Omega_\Lambda = 0.74$ and $\Omega_m = 0.26$ (dashed line), and for our model (solid line) with the values of the best-fit coming from SNIa+BAO+CMB, $\Omega_m = 0.27$ and $n = -0.10$. Three cosmological phases observed.




FIG. 5: Effective equation of state as a function of z for our model with the values of the best-fit coming from SNIa+BAO+CMB, $\Omega_m = 0.27$ and $n = -0.10$.

the theory $f(L_T) = L_T - \alpha(-L_T)^{-n}$ with the aim of reproducing the recently detected acceleration of the universe without resorting to dark energy. We have here performed observational viability tests for this theory by using the most recent SN Ia data, and combined them with the information coming from BAO peak and CMB shift parameter in order to find constraints in the $n - \Omega_m$ plane. At 68.3% c.l. we found that the values lie in the ranges $n \in [-0.23, 0.03]$ and $\Omega_m \in [0.25, 0.29]$. The values for Ω_m are consistent with recent estimations obtained by other authors (see, e.g., [39]). The model with the best-fit values minimizing the χ^2 that combines SNIa+BAO+CMB data ($n = -0.10$ and $\Omega_m = 0.27$) exhibits the last three phases of cosmological evolution: radiation era, matter era and late acceleration; this last

stage having started at $z_{acc} \simeq 0.74$.

Acknowledgments

G.R.B. is supported by a CONICET graduate scholarship. G.R.B thanks Franco Fiorini for his encouraging at

the early stages of the work. Both authors are grateful to Diego Travieso, Gastón Giribet and Susana Landau for helpful discussions. This work was partially supported by CONICET.

-
- [1] Perlmutter S. et al., Bull. Am. Astron. Soc. **29**, (1997) 1351; Astrophys. J. **517**, (1999) 565; Riess A. G. et al., Astrophys. J. **116**, (1998) 1009; Astrophys. J. **607** (2004) 665.
 - [2] Kowalski M. et al., Astrophys. J. **686**, (2008) 749.
 - [3] Sahni V. and Starobinsky A. A., Int. J. Mod. Phys. **D9**, (2000) 373.
 - [4] Padmanabhan T., Phys. Rep. **380**, (2003) 235.
 - [5] Frieman J., Turner M. S. and Huterer D., *Dark Energy and the Accelerating Universe*, arXiv:0803.0982.
 - [6] Buchdahl H. A., Mon. Not. R. Astron. Soc. **150**, (1970) 1.
 - [7] Starobinsky A. A., Phys. Lett. **B91**, (1980) 99.
 - [8] Kerner R., Gen. Relat. Gravit. **14**, (1982) 453.
 - [9] Barrow J. D. and Ottewill A. C., J. Phys. A: Math. Gen. **16**, (1983) 2757.
 - [10] Barrow J. D. and Cotsakis S., Phys. Lett. **B214**, (1988) 515.
 - [11] Carroll S. M. et al., Phys. Rev. **D70**, (2004) 043528.
 - [12] Starobinsky A. A., JETP Lett. **86**, (2007) 157.
 - [13] Hu W. and Sawicki I., Phys. Rev. **D76**, (2007) 064004.
 - [14] Nojiri S. and Odintsov S. D., *Dark energy, inflation and dark matter from modify $f(R)$ gravity*, arXiv:0807.0685.
 - [15] Nojiri S. and Odintsov S. D., Int. J. Geom. Meth. Mod. Phys. **4**, (2007) 115.
 - [16] Capozziello S. and Francaviglia M., Gen. Relat. Gravit. **40**, (2008) 357.
 - [17] Sotiriou T. and Faraoni V., *$f(R)$ theories of gravity*, arXiv:0805.1726.
 - [18] Chiba T., Phys. Lett. **B575**, (2003) 1.
 - [19] Olmo G., Phys. Rev. Lett. **95**, (2005) 261102.
 - [20] Dolgov A. and Kawasaki M., Phys. Lett. **B573**, (2003) 1.
 - [21] Amendola L. et al., Phys. Rev. Lett. **98**, (2007) 131302.
 - [22] Amendola L. et al., Phys. Rev. **D75**, (2007) 083504.
 - [23] Amarzguioui M. et al., Astron. and Astrophys. **454**, (2006) 707.
 - [24] Fay S., Tavakol R. and Tsujikawa S., Phys. Rev. **D75**, (2007) 063509.
 - [25] Santos J. et al., Phys. Lett. **B669**, (2008) 14.
 - [26] Ferraro R. and Fiorini F., Phys. Rev. **D75**, (2007) 084031.
 - [27] Ferraro R. and Fiorini F., Phys. Rev. **D78**, (2008) 124019.
 - [28] Einstein A., Sitzungsber. Preuss. Akad. Wiss. (1928) 217; ibid., (1930) 401; Einstein A., Math. Annal. **102**, (1930) 685.
 - [29] Hayashi K. and Shirafuji T., Phys. Rev. **D19**, (1979) 3524, Addendum-ibid. **D24**, (1982) 3312.
 - [30] Weitzenböck R., *Invarianten Theorie*, (Nordhoff, Groningen, 1923).
 - [31] Maluf J. W., J. Math. Phys. **35**, (1994) 335.
 - [32] Arcos H. and Pereira J., Int. J. Mod. Phys. **D13**, (2004) 2193.
 - [33] Dvali G. and Turner M. S., *Dark Energy as a Modification of the Friedmann Equation*, ArXiv:astro-ph/0301510.
 - [34] Fairbairn M. and Goobar A., Phys. Lett. **B642**, (2006) 432.
 - [35] Eisenstein D. et al., Astrophys. J. **633**, (2005) 560.
 - [36] Komatsu E. et al., Astrophys. J. Suppl. **180**, (2009) 330.
 - [37] Melchiorri A. et al., Phys. Rev. **D76**, (2007) 041301(R).
 - [38] Caldwell R. R., Phys. Lett. **B545**, (2002) 23.
 - [39] Spergel D. N. et al., Astrophys. J. Suppl., **148**, (2003) 175; Astrophys. J. Suppl., **170**, (2007) 377.